

ON α -EMBEDDED SETS AND EXTENSION OF MAPPINGS

OLENA KARLOVA

ABSTRACT. We introduce and study α -embedded sets and apply them to generalize the Kuratowski Extension Theorem.

1. INTRODUCTION

A subset A of a topological space X is called *functionally open* (*functionally closed*) if there exists a continuous function $f : X \rightarrow [0, 1]$ such that $A = f^{-1}((0, 1])$ ($A = f^{-1}(0)$).

Let $\mathcal{G}_0^*(X)$ and $\mathcal{F}_0^*(X)$ be the collections of all functionally open and functionally closed subsets of a topological space X , respectively. Assume that the classes $\mathcal{G}_\xi^*(X)$ and $\mathcal{F}_\xi^*(X)$ are defined for all $\xi < \alpha$, where $0 < \alpha < \omega_1$. Then, if α is odd, the class $\mathcal{G}_\alpha^*(X)$ ($\mathcal{F}_\alpha^*(X)$) consists of all countable intersections (unions) of sets of lower classes, and, if α is even, the class $\mathcal{G}_\alpha^*(X)$ ($\mathcal{F}_\alpha^*(X)$) consists of all countable unions (intersections) of sets of lower classes. The classes $\mathcal{F}_\alpha^*(X)$ for odd α and $\mathcal{G}_\alpha^*(X)$ for even α are said to be *functionally additive*, and the classes $\mathcal{F}_\alpha^*(X)$ for even α and $\mathcal{G}_\alpha^*(X)$ for odd α are called *functionally multiplicative*. If a set belongs to the α 'th functionally additive and to the α 'th functionally multiplicative class simultaneously, then it is called *functionally ambiguous of the α 'th class*. For every $0 \leq \alpha < \omega_1$ let

$$\mathcal{B}_\alpha^*(X) = \mathcal{F}_\alpha^*(X) \cup \mathcal{G}_\alpha^*(X)$$

and let

$$\mathcal{B}^*(X) = \bigcup_{0 \leq \alpha < \omega_1} \mathcal{B}_\alpha^*(X).$$

If $A \in \mathcal{B}^*(X)$, then A is said to be a *functionally measurable set*.

If P is a property of mappings, then by $P(X, Y)$ we denote the collection of all mappings $f : X \rightarrow Y$ with the property P . Let $P(X)$ ($P^*(X)$) be the collection of all real-valued (bounded) mappings on X with a property P .

By the symbol C we denote, as usually, the property of continuity.

Let $K_0(X, Y) = C(X, Y)$. For an ordinal $0 < \alpha < \omega_1$ we say that a mapping $f : X \rightarrow Y$ belongs to the α 'th functional Lebesgue class, $f \in K_\alpha(X, Y)$, if the preimage $f^{-1}(V)$ of an arbitrary open set $V \subseteq Y$ is of the α 'th functionally additive class in X .

A subspace E of X is *P-embedded* (*P*-embedded*) in X if every (bounded) function $f \in P(E)$ can be extended to a (bounded) function $g \in P(X)$.

A subset E of X is said to be *z-embedded* in X if every functionally closed set in E is the restriction of a functionally closed set in X to E . It is well-known that

$$E - C\text{-embedded} \Rightarrow E - C^*\text{-embedded} \Rightarrow E - z\text{-embedded}.$$

Recall that sets A and B are *completely separated* in X if there exists a continuous function $f : X \rightarrow [0, 1]$ such that $A \subseteq f^{-1}(0)$ and $B \subseteq f^{-1}(1)$.

The following theorem was proved in [2, Corollary 3.6].

Theorem 1.1 (Blair-Hager). *A subset E of a topological space X is C -embedded in X if and only if E is z -embedded in X and E is completely separated from every functionally closed set in X disjoint from E .*

It is natural to consider P - and P^* -embedded sets if $P = K_\alpha$ for $\alpha > 0$. In connection with this we introduce and study a class of α -embedded sets which coincides with the class of z -embedded sets when $\alpha = 0$. In Section 3 we generalize the notion of completely separated sets to α -separated sets. Section 4 deals with ambiguously α -embedded sets which play the important role in the extension of bounded K_α -functions. In the fifth section we prove an analog of the Tietze-Uryhson Extension Theorem for K_α -functions. Section 6 concerns the question when K_1 -embedded sets coincide with K_1^* -embedded sets. The seventh section presents a generalization of the Kuratowski Theorem [11, p. 445] on extension of K_α -mappings with values in Polish spaces.

2. α -EMBEDDED SETS

Let $0 \leq \alpha < \omega_1$. A subset E of a topological space X is *α -embedded* in X if for any set A of the α 'th functionally additive (multiplicative) class in E there is a set B of the α 'th functionally additive (multiplicative) class in X such that $A = B \cap E$.

Proposition 2.1. *Let X be a topological space, $0 \leq \alpha < \omega_1$ and let $E \subseteq X$ be an α -embedded set of the α 'th functionally additive (multiplicative) class in X . Then every set of the α 'th functionally additive (multiplicative) class in E belongs to the α 'th functionally additive (multiplicative) class in X .*

Proof. For a set C of the α 'th functionally additive (multiplicative) class in E we choose a set B of the α 'th functionally additive (multiplicative) class in X such that $C = B \cap E$. Then C belongs to the α 'th functionally additive (multiplicative) class in X as the intersection of two sets of the same class. $\square$

Proposition 2.2. *Let X be a topological space, $E \subseteq X$ and*

- (i) *X is perfectly normal, or*
- (ii) *X is completely regular and E is its Lindelöf subset, or*
- (iii) *E is a functionally open subset of X , or*
- (iv) *X is a normal space and E is its F_σ -subset,*

then E is 0-embedded in X .

Proof. Let G be a functionally open set in E .

(i). Choose an open set U in X such that $G = E \cap U$. Then U is functionally open in X by Vedenissoff's theorem [5, p. 45].

(ii). Let U be an open set in X such that $G = E \cap U$. Since X is completely regular, $U = \bigcup_{s \in S} U_s$, where U_s is a functionally open set in X for each $s \in S$.

Notice that G is Lindelöf, provided G is F_σ in the Lindelöf space E [5, p. 192]. Then there exists a countable set $S_0 \subseteq S$ such that $G \subseteq \bigcup_{s \in S_0} U_s$. Let $V = \bigcup_{s \in S_0} U_s$.

Then V is functionally open in X and $V \cap E = G$.

(iii). Consider continuous functions $\varphi : E \rightarrow [0, 1]$ and $\psi : X \rightarrow [0, 1]$ such that $G = \varphi^{-1}((0, 1])$ and $E = \psi^{-1}((0, 1])$. For each $x \in X$ we set

$$f(x) = \begin{cases} \varphi(x) \cdot \psi(x), & x \in E, \\ 0, & x \in X \setminus E. \end{cases}$$

Since $\varphi(x) \cdot \psi(x) = 0$ on $\overline{E} \setminus E$, $f : X \rightarrow [0, 1]$ is continuous. Moreover, $G = f^{-1}((0, 1])$. Hence, the set G is functionally open in X .

(iv). Let $\tilde{G}$ be an open set in X such that $G = \tilde{G} \cap E$. Since G is functionally open in E , G is F_σ in E . Consequently, G is F_σ in X , provided E is F_σ in X . Since X is normal, for every $n \in \mathbb{N}$ there exists a continuous function $f_n : X \rightarrow [0, 1]$ such that $f_n(x) = 1$ if $x \in F_n$ and $f_n(x) = 0$ if $x \in X \setminus \tilde{G}$. Then the set $V = \bigcup_{n=1}^{\infty} f_n^{-1}((0, 1])$ is functionally open in X and $V \cap E = G$. $\square$

Examples 2.3 and 2.4 show that none of the conditions (i)–(iv) on X and E in Proposition 2.2 can be weakened.

Recall that a topological space X is said to be *perfect* if every its closed subset is G_δ in X .

Example 2.3. *There exist a perfect completely regular space X and its functionally closed subspace E which is not α -embedded in X for every $0 \leq \alpha < \omega_1$.*

Consequently, there is a bounded continuous function on E which cannot be extended to a $\mathcal{K}_\alpha$ -function for every α .

Proof. Let X be the Niemytski plane [5, p. 22], i.e. $X = \mathbb{R} \times [0, +\infty)$, where a base of neighborhoods of $(x, y) \in X$ with $y > 0$ form open balls with the center in (x, y) , and a base of neighborhoods of $(x, 0)$ form the sets $U \cup \{(x, 0)\}$, where U is an open ball which tangent to $\mathbb{R} \times \{0\}$ in the point $(x, 0)$. It is well-known that the space X is perfect and completely regular, but is not normal.

Denote $E = \mathbb{R} \times \{0\}$. Since the function $f : X \rightarrow \mathbb{R}$, $f(x, y) = y$, is continuous and $E = f^{-1}(0)$, the set E is functionally closed in X .

Notice that every function $f : E \rightarrow \mathbb{R}$ is continuous. Therefore, $|\mathcal{B}_\alpha^*(E)| = 2^{2^{\omega_0}}$ for every $0 \leq \alpha < \omega_1$. On the other hand, $|\mathcal{B}_\alpha^*(X)| = 2^{\omega_0}$ for every $0 \leq \alpha < \omega_1$, provided the space X is separable. Hence, for every $0 \leq \alpha < \omega_1$ there exists a set $A \in \mathcal{B}_\alpha^*(E)$ which can not be extended to a set $B \in \mathcal{B}_\alpha^*(X)$.

Observe that a function $f : E \rightarrow [0, 1]$, such that $f = 1$ on A and $f = 0$ on $E \setminus A$, is continuous on E . But there is no K_α -function $f : X \rightarrow [0, 1]$ such that $g|_E = f$, since otherwise the set $B = g^{-1}(1)$ would be an extension of A . $\square$

Example 2.4. *There exist a compact Hausdorff space X and its open subspace E which is not α -embedded in X for every $0 \leq \alpha < \omega_1$.*

Proof. Let $X = D \cup \{\infty\}$ be the Alexandroff compactification of an uncountable discrete space D [5, p. 169] i $E = D$. Fix $0 \leq \alpha < \omega_1$ and choose an arbitrary uncountable set $A \subseteq E$ with uncountable complement $X \setminus A$. Evidently, A is functionally closed in E . Assume that there is a set B of the α 'th functionally multiplicative class in X such that $A = B \cap E$. Clearly, $B = A \cup \{\infty\}$. Moreover, there exists a function $f : X \rightarrow \mathbb{R}$ of the α 'th Baire class such that $B = f^{-1}(0)$ [9, Lemma 2.1]. But every continuous function on X , and consequently every Baire function of the class α on X satisfies the equality $f(x) = f(\infty)$ for all but countably many points $x \in X$, which implies a contradiction. $\square$

Proposition 2.5. *Let $0 \leq \alpha \leq \beta < \omega_1$ and let X be a topological space. Then every α -embedded subset of X is β -embedded.*

Proof. Let E be an α -embedded subset of X . If $\beta = \alpha$, the assertion of the proposition is obvious. Suppose the assertion is true for all $\alpha \leq \beta < \xi$ and let A be a set of the ξ 'th functionally additive class in E . Then there exists a sequence of sets A_n of functionally multiplicative classes $< \xi$ in E such that $A = \bigcup_{n=1}^{\infty} A_n$. According to the assumption, for every $n \in \mathbb{N}$ there is a set B_n of a functionally multiplicative class $< \xi$ in X such that $A_n = B_n \cap E$. Then the set $B = \bigcup_{n=1}^{\infty} B_n$ belongs to the ξ 'th functionally additive class in X and $A = B \cap E$. $\square$

The inverse proposition is not true, as the following result shows.

Theorem 2.6. *There exist a completely regular space X and its 1-embedded subspace $E \subseteq X$ which is not 0-embedded in X .*

Proof. Let $X_0 = [0, 1]$, $X_s = \mathbb{N}$ for every $s \in (0, 1]$, $Y = \prod_{s \in (0, 1]} X_s$ and

$$X = [0, 1] \times Y = \prod_{s \in [0, 1]} X_s.$$

Then X is completely regular as a product of completely regular spaces X_s . Let

$$A_1 = (0, 1] \quad \text{and} \quad A_2 = \{0\}.$$

For $i = 1, 2$ we consider the set

$$F_i = \bigcap_{n \neq i} \{y = (y_s)_{s \in (0, 1]} \in Y : |\{s \in (0, 1] : y_s = n\}| \leq 1\}.$$

Obviously, $F_1 \cap F_2 = \emptyset$ and the sets F_1 and F_2 are closed in Y .

Let

$$B_1 = A_1 \times F_1, \quad B_2 = A_2 \times F_2 \quad \text{and} \quad E = B_1 \cup B_2.$$

It is easy to see that the sets B_1 and B_2 are closed in E , and consequently they are functionally clopen in E .

Claim 1. The set B_i is 0-embedded in X for every $i = 1, 2$.

Proof. Let C be a functionally open set in B_1 .

Let us consider the set

$$H = \{x = (x_s)_{s \in [0,1]} \in X : |\{s \in [0,1] : x_s \neq 1\}| \leq \aleph_0\}.$$

Then the set $[0, 1] \times F_i$ is closed in H for every $i = 1, 2$. Since H is the Σ -product of the family $(X_s)_{s \in [0,1]}$ (see [5, p. 118]), according to [10] the space H is normal. Consequently, $[0, 1] \times F_i$ is normal as closed subspace of normal space for every $i = 1, 2$. Clearly, B_1 is functionally open in $[0, 1] \times F_1$. Hence, B_1 is 0-embedded in $[0, 1] \times F_1$ according to Proposition 2.2(iii). Then C is functionally open in $[0, 1] \times F_1$ by Proposition 2.1. Notice that the set $[0, 1] \times F_1$ is 0-embedded in H by Propositions 2.2 (iv). Hence, there exists a functionally open set C' in H such that $C' \cap ([0, 1] \times F_1) = C$. It follows from [3] that H is 0-embedded in X . Then there exists a functionally open set C'' in X such that $C'' \cap H = C'$. Evidently, $C'' \cap B_1 = C$. Therefore, the set B_1 is 0-embedded in X .

Analogously, it can be shown that the set B_2 is 0-embedded in X , using the fact that B_2 is 0-embedded in $[0, 1] \times F_2$ according to Proposition 2.2(iv).

Claim 2. The set E is not 0-embedded in X .

Proof. Assuming the contrary, we choose a functionally closed set D in X such that $D \cap E = B_1$. Then $D = f^{-1}(0)$ for some continuous function $f : X \rightarrow [0, 1]$. It follows from [5, p. 117] that there exists a countable set $S = \{0\} \cup T$, where $T \subseteq (0, 1]$, such that for any $x = (x_s)_{s \in [0,1]}$ and $y = (y_s)_{s \in [0,1]}$ of X the equality $x|_S = y|_S$ implies $f(x) = f(y)$. Let $y_0 \in Y$ be such that $y_0|_T$ is a sequence of different natural numbers which are not equal to 1 or 2. We choose $y_1 \in F_1$ and $y_2 \in F_2$ such that $y_0|_T = y_1|_T = y_2|_T$. Then

$$f(a, y_0) = f(a, y_1) = f(a, y_2)$$

for all $a \in [0, 1]$. We notice that $f(0, y_1) = 0$. Therefore, $f(0, y_0) = 0$. But $f(a, y_2) > 0$ for all $a \in A_2$. Then $f(a, y_0) > 0$ for all $a \in A_2$. Hence, $A_1 = (f^{y_0})^{-1}(0)$, where $f^{y_0}(a) = f(a, y_0)$ for all $a \in [0, 1]$, and f^{y_0} is continuous. Thus, the set $A_1 = (0, 1]$ is closed in $[0, 1]$, which implies a contradiction.

Claim 3. The set E is 1-embedded in X .

Proof. Let C be a functionally G_δ -set in E . We put

$$E_1 = A_1 \times Y, \quad E_2 = A_2 \times Y.$$

Then the set E_1 is functionally open in X and the set E_2 is functionally closed in X . For $i = 1, 2$ let $C_i = C \cap B_i$. Since for every $i = 1, 2$ the set C_i is functionally

G_δ in the 0-embedded in X set B_i , by Proposition 2.5 there exists a functionally G_δ -set $\tilde{C}_i$ in X such that $\tilde{C}_i \cap B_i = C_i$. Let

$$\tilde{C} = (\tilde{C}_1 \cap E_1) \cup (\tilde{C}_2 \cap E_2).$$

Then $\tilde{C}$ is functionally G_δ in X and $\tilde{C} \cap E = C$. $\square$

3. α -SEPARATED SETS AND α -SEPARATED SPACES

Let $0 \leq \alpha < \omega_1$. Subsets A and B of a topological space X are said to be α -separated if there exists a function $f \in K_\alpha(X)$ such that

$$A \subseteq f^{-1}(0) \quad \text{and} \quad B \subseteq f^{-1}(1).$$

Remark that 0-separated sets are also called *completely separated* [5, p. 42].

Lemma 3.1 (Lemma 2.1 [8]). *Let X be a topological space, $\alpha > 0$ and let $A \subseteq X$ be a subset of the α 'th functionally additive class. Then there exists a sequence $(A_n)_{n=1}^\infty$ such that each A_n is functionally ambiguous of the class α in X , $A_n \cap A_m = \emptyset$ for $n \neq m$ and $A = \bigcup_{n=1}^\infty A_n$.*

Proof. Since A belongs to the α 'th functionally additive class, $A = \bigcup_{n=1}^\infty B_n$, where each B_n belongs to the functionally multiplicative class $< \alpha$ in X . Therefore, each B_n is functionally ambiguous of the class α . Let $A_1 = B_1$ and $A_n = B_n \setminus \bigcup_{k < n} B_k$ for $n > 1$. Then $(A_n)_{n=1}^\infty$ is the required sequence. $\square$

Lemma 3.2 (Lemma 2.2 [8]). *Let X be a topological space, $\alpha \geq 0$ and let A_n belongs to the α 'th functionally additive class in X for every $n \in \mathbb{N}$ with $X = \bigcup_{n=1}^\infty A_n$. Then there exists a sequence $(B_n)_{n=1}^\infty$ of mutually disjoint functionally ambiguous sets of the class α in X such that $B_n \subseteq A_n$ and $X = \bigcup_{n=1}^\infty B_n$.*

Proof. It follows from Lemma 3.1 that for every $n \in \mathbb{N}$ there exists a sequence $(F_{n,m})_{m=1}^\infty$ such that each $F_{n,m}$ is functionally ambiguous of the class α in X , $F_{n,m} \cap F_{n,k} = \emptyset$ for $m \neq k$ and $A_n = \bigcup_{m=1}^\infty F_{n,m}$. Let $k : \mathbb{N}^2 \rightarrow \mathbb{N}$ be a bijection. Set

$$C_{n,m} = F_{n,m} \setminus \bigcup_{k(p,s) < k(n,m)} F_{p,s}.$$

Evidently, $\bigcup_{n,m=1}^\infty C_{n,m} = X$. Let $B_n = \bigcup_{m=1}^\infty C_{n,m}$. Then $\bigcup_{n=1}^\infty B_n = \bigcup_{n=1}^\infty A_n = X$ and $B_n \subseteq \bigcup_{m=1}^\infty F_{n,m} = A_n$. Notice that each $C_{n,m}$ is functionally ambiguous of the class α . Therefore, B_n belongs to the functionally additive class α for every n . Moreover, $B_n \cap B_m = \emptyset$ for $n \neq m$. Since $X \setminus B_n = \bigcup_{k \neq n} B_k$, B_n is functionally ambiguous of the class α . $\square$

Lemma 3.3. *Let $0 \leq \alpha < \omega_1$ and let A be a subset of the α 'th functionally multiplicative class of a topological space X . Then there exists a function $f \in K_\alpha^*(X)$ such that $A = f^{-1}(0)$.*

Proof. For $\alpha = 0$ the lemma implies from the definition of a functionally closed set.

Let $\alpha > 0$. Since the set $B = X \setminus A$ is of the α 'th functionally additive class, there exists a sequence of functionally ambiguous sets B_n of the α 'th class in X such that $B = \bigcup_{n=1}^{\infty} B_n$ and $B_n \cap B_m = \emptyset$ for all $n \neq m$ by Lemma 3.1. Define a function $f : X \rightarrow [0, 1]$,

$$f(x) = \begin{cases} 0, & \text{if } x \in A, \\ \frac{1}{n}, & \text{if } x \in B_n. \end{cases}$$

Take an arbitrary open set $V \subseteq [0, 1]$. If $0 \notin V$ then $f^{-1}(V)$ is of the α 'th functionally additive class as a union of at most countably many sets B_n . If $0 \in V$ then there exists such a number N that $\frac{1}{n} \in V$ for all $n > N$. Then the set $X \setminus f^{-1}(V) = \bigcup_{n=1}^N B_n$ belongs to the α 'th functionally multiplicative class. Hence, $f^{-1}(V)$ is of the α 'th functionally additive class in X . Therefore, $f \in K_\alpha^*(X)$. $\square$

Proposition 3.4. *Let $0 \leq \alpha < \omega_1$ and let X be a topological space. Then any two disjoint sets A and B of the α 'th functionally multiplicative class in X are α -separated.*

Proof. By Lemma 3.3 we choose functions $f_1, f_2 \in K_\alpha(X)$ such that $A = f_1^{-1}(0)$ and $B = f_2^{-1}(0)$. For all $x \in X$ let

$$f(x) = \frac{f_1(x)}{f_1(x) + f_2(x)}.$$

It is easy to see that $f \in K_\alpha(X)$, $f(x) = 0$ on A and $f(x) = 1$ on B . $\square$

Let $0 \leq \alpha < \omega_1$. A topological space X is α -separated if any two disjoint sets $A, B \subseteq X$ of the α 'th multiplicative class in X are α -separated. It follows from Urysohn's Lemma [5, p. 41] that a topological space is 0-separated if and only if it is normal. Proposition 3.4 implies that every perfectly normal space is α -separated for each $\alpha \geq 0$. It is naturally to ask whether there is an α -separated space for $\alpha \geq 1$ which is not perfectly normal.

Example 3.5. *There exists a completely regular 1-separated space which is not perfectly normal.*

Proof. Let $D = D(\mathfrak{m})$ be a discrete space of the cardinality $\mathfrak{m}$, where $\mathfrak{m}$ is a measurable cardinal number [6, 12.1]. According to [6, 12.2], D is not a realcompact space. Let $X = vD$ be a Hewitt realcompactification of D [5, p. 218]. Then X is an extremally disconnected P -space, which is not discrete [6, 12H]. Thus, there exists a point $x \in X$ such that the set $\{x\}$ is not open. Then $\{x\}$, being a closed

set, is not a G_δ -set, since X is a P -space (i.e. a space in which every G_δ -subset is open). Therefore, the space X is not perfect.

If A and B are disjoint G_δ -subsets of X , then A and B are open in X . Notice that in an extremally disconnected space any two disjoint open sets are completely separated [6, 1H]. Consequently, A and B are 1-separated, since every continuous function belongs to the first Lebesgue class. $\square$

Clearly, every ambiguous set A of the class 0 in a topological space (i.e., every clopen set) is a functionally ambiguous set of the class 0. If A is an ambiguous set of the first class, i.e. A is an F_σ - and a G_δ -set, then A need not be a functionally F_σ - or a functionally G_δ -set. Indeed, let X be the Niemytski plane, E be a set which is not of the G_δ -type in $\mathbb{R}$ and let $A = E \times \{0\}$ be a subspace of X . Then A is closed and consequently G_δ -subset of X , since the Niemytski plane is a perfect space. Assume that A is a functionally F_σ -set in X . Then $A = \bigcup_{n=1}^{\infty} A_n$, where A_n is a functionally closed subset of X for every $n \in \mathbb{N}$. According to [13, Theorem 5.1], a closed subset F of X is a functionally closed set in X if and only if the set $\{x \in \mathbb{R} : (x, 0) \in F\}$ is a G_δ -set in $\mathbb{R}$. It follows that for every $n \in \mathbb{N}$ the set A_n is a G_δ -subset of $\mathbb{R}$, which implies a contradiction.

Theorem 3.6. *Let $0 \leq \alpha < \omega_1$ and let X be an α -separated space.*

- (1) *Every ambiguous set $A \subseteq X$ of the class α is functionally ambiguous of the class α .*
- (2) *For any disjoint sets A and B of the $(\alpha + 1)$ 'th additive class in X there exists a set C of the $(\alpha + 1)$ 'th functionally multiplicative class such that*

$$A \subseteq C \subseteq X \setminus B.$$

- (3) *Every ambiguous set A of the $(\alpha + 1)$ 'th class in X is a functionally ambiguous set of the $(\alpha + 1)$ 'th class.*
- (4) *Any set of the α 'th multiplicative class in X is α -embedded.*

Proof. (1) Since the set $B = X \setminus A$ belongs to the α 'th multiplicative class in X , there exists a function $f \in K_\alpha(X)$ such that $A \subseteq f^{-1}(0)$ and $B \subseteq f^{-1}(1)$. Then $A = f^{-1}(0)$ and $B = f^{-1}(1)$. Hence, the sets A and B are of the α 'th functionally multiplicative class. Consequently, A is a functionally ambiguous set of the class α .

(2) Choose two sequences $(A_n)_{n=1}^{\infty}$ and $(B_n)_{n=1}^{\infty}$, where A_n and B_n belong to the α 'th multiplicative class in X for every $n \in \mathbb{N}$, such that $A = \bigcup_{n=1}^{\infty} A_n$ and $B = \bigcup_{n=1}^{\infty} B_n$. Since X is α -separated, for every $n, m \in \mathbb{N}$ there exists a function $f_{n,m} \in K_\alpha(X)$ such that $A_n \subseteq f_{n,m}^{-1}(1)$ and $B_m \subseteq f_{n,m}^{-1}(0)$. Set

$$C = \bigcap_{n=1}^{\infty} \bigcup_{m=1}^{\infty} f_{n,m}^{-1}((0, 1]).$$

Then the set C is of the $(\alpha + 1)$ 'th functionally multiplicative class in X and $A \subseteq C \subseteq X \setminus B$.

(3) Let $A \subseteq X$ be an ambiguous set of the $(\alpha + 1)$ 'th class. Denote $B = X \setminus A$. Since A and B are disjoint sets of the $(\alpha + 1)$ 'th additive class in X , according to (2) there exists a set $C \subseteq X$ of the $(\alpha + 1)$ 'th functionally multiplicative class such that $A \subseteq C \subseteq X \setminus B$. It follows that $A = C$, consequently A is of the $(\alpha + 1)$ 'th functionally multiplicative class. Analogously, it can be shown that B is also of the $(\alpha + 1)$ 'th functionally multiplicative class. Therefore, A is a functionally ambiguous set of the $(\alpha + 1)$ 'th class.

(4) If $\alpha = 0$ then X is a normal space. Therefore, any closed set F in X is 0-embedded by Proposition 2.2.

Let $\alpha > 0$ and let $E \subseteq X$ be a set of the α 'th multiplicative class in X . Choose any set A of the α 'th functionally multiplicative class in E . Since the set $E \setminus A$ belongs to the α 'th functionally additive class in E , there exists a sequence of sets B_n of the α 'th functionally multiplicative class in E such that $E \setminus A = \bigcup_{n=1}^{\infty} B_n$. Then for every $n \in \mathbb{N}$ the sets A and B_n are disjoint and belong to the α 'th multiplicative class in X . Since X is α -separated, we can choose a function $f_n \in K_\alpha(X)$ such that $A \subseteq f_n^{-1}(0)$ and $B_n \subseteq f_n^{-1}(1)$. Let $\tilde{A} = \bigcap_{n=1}^{\infty} f_n^{-1}(0)$. Then the set $\tilde{A}$ belongs to the α 'th functionally multiplicative class in X and $\tilde{A} \cap E = A$. $\square$

Proposition 3.7. *A topological space X is normal if and only if every its closed subset is 0-embedded.*

Proof. We only need to prove the sufficiency. Let A and B be disjoint closed subsets of X . Then A is a functionally closed subset of $E = A \cup B$. Since E is closed in X , E is a 0-embedded set. Therefore, there is a functionally closed set $\tilde{A}$ in X such that $A = E \cap \tilde{A}$. Then B is a functionally closed subset of the closed set $D = \tilde{A} \cup B$. Since D is 0-embedded in X , there exists a functionally closed set $\tilde{B}$ in X such that $B = D \cap \tilde{B}$. It is easy to check that $\tilde{A} \cap \tilde{B} = \emptyset$. If $f : X \rightarrow [0, 1]$ be a continuous function such that $\tilde{A} = f^{-1}(0)$ and $\tilde{B} = f^{-1}(1)$, then the sets $U = f^{-1}([0, 1/2))$ and $V = f^{-1}((1/2, 1])$ are disjoint and open in X , $A \subseteq U$ and $B \subseteq V$. Hence, X is a normal space. $\square$

An analog of the previous proposition takes place for hereditarily α -separated spaces. We say that a topological space X is *hereditarily α -separated* if every its subspace is α -separated.

Proposition 3.8. *Let $0 \leq \alpha < \omega_1$ and let X be a hereditarily α -separated space. If every subset of the $(\alpha + 1)$ 'th multiplicative class in X is $(\alpha + 1)$ -embedded, then X is $(\alpha + 1)$ -separated.*

Proof. Let $A, B \subseteq X$ be disjoint sets of the $(\alpha + 1)$ 'th multiplicative class. Then A is ambiguous of the class $(\alpha + 1)$ in $E = A \cup B$. Since E belongs to the $(\alpha + 1)$ 'th multiplicative class in X , E is $(\alpha + 1)$ -embedded. Moreover, E is α -separated as

a subspace of the hereditarily α -separated space X . According to Theorem 3.6(3) A is functionally ambiguous of the $(\alpha + 1)$ 'th class in E . Therefore, there is a set $\tilde{A}$ of the $(\alpha + 1)$ 'th functionally multiplicative class in X such that $A = E \cap \tilde{A}$. Then B is a functionally ambiguous subset of the class $(\alpha + 1)$ in $D = \tilde{A} \cup B$. Since D belongs to the $(\alpha + 1)$ 'th multiplicative class in X , D is $(\alpha + 1)$ -embedded. Therefore, there exists a set $\tilde{B}$ of the $(\alpha + 1)$ 'th functionally multiplicative class in X such that $B = D \cap \tilde{B}$. It is easy to check that $\tilde{A} \cap \tilde{B} = \emptyset$. Hence, the sets $\tilde{A}$ and $\tilde{B}$ are $(\alpha + 1)$ -separated by Proposition 3.4. Then A and B are $(\alpha + 1)$ -separated too. $\square$

Remark that the Alexandroff compactification of the real line $\mathbb{R}$ endowed with the discrete topology is a hereditarily normal space which is not 1-separated.

We give some examples below of α -separated subsets of a completely regular space.

Proposition 3.9. *Let X be a completely regular space and $A, B \subseteq X$ are disjoint sets. Then*

- (a) *if A and B are Lindelöf G_δ -sets, then they are 1-separated;*
- (b) *if A is a Lindelöf hereditarily Baire space and B is a functionally G_δ -set, then A and B are 1-separated;*
- (c) *if A is Lindelöf and B is an F_σ -set, then A and B are 2-separated.*

Proof. (a). Let $A = \bigcap_{n=1}^{\infty} U_n$, where U_n is an open set in X for every $n \in \mathbb{N}$. Since X is completely regular, $U_n = \bigcup_{s \in S_n} U_{s,n}$ for every $n \in \mathbb{N}$, where all the sets $U_{s,n}$ are functionally open in X . Then for every $n \in \mathbb{N}$ there is a countable set $S_{n,0} \subseteq S_n$ such that $A \subseteq \bigcup_{s \in S_{n,0}} U_{s,n}$, since A is Lindelöf. Let $V_n = \bigcup_{s \in S_{n,0}} U_{s,n}$, $n \in \mathbb{N}$. Obviously, every V_n is a functionally open set and $A = \bigcap_{n=1}^{\infty} V_n$. Hence, A is a functionally G_δ -subset of X . Analogously, B is also a functionally G_δ -set. Therefore, the sets A and B are 1-separated by Proposition 3.4.

(b). According to [7, Proposition 12] there is a functionally G_δ -set C in X such that $A \subseteq C \subseteq X \setminus B$. Taking a function $f \in K_1(X)$ such that $C = f^{-1}(0)$ and $B = f^{-1}(1)$, we obtain that A and B are 1-separated.

(c). Let $X \setminus B = \bigcap_{n=1}^{\infty} U_n$, where $(U_n)_{n=1}^{\infty}$ is a sequence of open subsets of X . Then $U_n = \bigcup_{s \in S_n} U_{s,n}$ for every $n \in \mathbb{N}$, where all the sets $U_{s,n}$ are functionally open in X . Since A is Lindelöf, $A \subseteq V_n = \bigcup_{s \in S_{n,0}} U_{s,n}$, where the set $S_{n,0}$ is countable for every $n \in \mathbb{N}$. Denote $C = \bigcap_{n=1}^{\infty} V_n$. Then C is a functionally G_δ -set in X and $A \subseteq C \subseteq X \setminus B$. Since C is a functionally ambiguous set of the second class, A and B are 2-separated. $\square$

The following example shows that the class of separation of sets A and B in Proposition 3.9(c) can not be made lower.

Example 3.10. *There exist a metrizable space X and its disjoint Lindelöf F_σ -subsets A and B , which are not 1-separated.*

Proof. Let $X = \mathbb{R}$, $A = \mathbb{Q}$ and B is a countable dense subsets of irrational numbers. Assume that A and B are 1-separated, i.e. there exist disjoint G_δ -sets C and D in $\mathbb{R}$ such that $A \subseteq C$ and $B \subseteq D$. Then $\overline{C} = \overline{D} = \mathbb{R}$, which implies a contradictions, since X is a Baire space. $\square$

4. AMBIGUOUSLY α -EMBEDDED SETS

Let $0 < \alpha < \omega_1$. A subset E of a topological space X is *ambiguously α -embedded in X* if for any functionally ambiguous set A of the class α in E there exists a functionally ambiguous set B of the class α in X such that $A = B \cap E$.

Proposition 4.1. *Let $0 < \alpha < \omega_1$ and let X be a topological space. Then every ambiguously α -embedded set E in X is α -embedded in X .*

Proof. Take a set $A \subseteq E$ of the α 'th functionally additive class in E . Then A can be written as $A = \bigcup_{n=1}^{\infty} A_n$, where A_n is a functionally ambiguous set of the class α in E for every $n \in \mathbb{N}$ by Lemma 3.1. Then there exists a sequence of functionally ambiguous sets B_n of the class α in X such that $A_n = B_n \cap E$ for every $n \in \mathbb{N}$. Let $B = \bigcup_{n=1}^{\infty} B_n$. Then the set B belongs to the α 'th functionally additive class in X and $B \cap E = A$. $\square$

We will need the following auxiliary fact.

Lemma 4.2 (Lemma 2.3 [8]). *Let $0 < \alpha < \omega_1$ and let X be a topological space. Then for any disjoint sets $A, B \subseteq X$ of the α 'th functionally multiplicative class in X there exists a functionally ambiguous set C of the class α in X such that $A \subseteq C \subseteq X \setminus B$.*

Proof. Lemma 3.2 implies that there are disjoint functionally ambiguous sets E_1 and E_2 of the class α such that $E_1 \subseteq X \setminus A$, $E_2 \subseteq X \setminus B$ and $X = E_1 \cup E_2$. It remains to put $C = E_2$. $\square$

Proposition 4.3. *Let $0 < \alpha < \omega_1$ and let X be a topological space. Then every α -embedded set E of the α 'th functionally multiplicative class in X is ambiguously α -embedded in X .*

Proof. Consider a functionally ambiguous set A of the class α in E . Then there exists a set B of the α 'th functionally multiplicative class in X such that $A = B \cap E$. Since E is of the α 'th functionally multiplicative class in X , the set A is also of the same class in X . Analogously, the set $E \setminus A$ belongs to the α 'th functionally multiplicative class in X . It follows from Lemma 4.2 that there exists a functionally ambiguous set C of the class α in X such that $A \subseteq C$ and

$C \cap (E \setminus A) = \emptyset$. Clearly, $C \cap E = A$. Hence, the set E is ambiguously α -embedded in X . $\square$

Example 4.4. *There exists a 0-embedded F_σ -set $E \subseteq \mathbb{R}$ which is not ambiguously 1-embedded.*

Proof. Let $E = \mathbb{Q}$. Obviously, E is a 0-embedded set. Consider any two disjoint A and B which are dense in E . Then A and B are simultaneously F_σ - and G_δ -sets in E . Assume that there exists an F_σ - and G_δ -set C in $\mathbb{R}$ such that $A = E \cap C$. Since $A \subseteq C$ and $B \subseteq \mathbb{R} \setminus C$, the sets C and $\mathbb{R} \setminus C$ are dense in $\mathbb{R}$. Moreover, the sets C and $\mathbb{R} \setminus C$ are G_δ in $\mathbb{R}$. It implies a contradiction, since $\mathbb{R}$ is a Baire space. $\square$

Example 4.5. *There exists a Borel non-measurable ambiguously 1-embedded subset of a perfectly normal compact space.*

Proof. Let X be the "two arrows" space (see [5, p. 212]), i.e. $X = X_0 \cup X_1$, where $X_0 = \{(x, 0) : x \in (0, 1]\}$ and $X_1 = \{(x, 1) : x \in [0, 1)\}$. The topology base on X is generated by the sets

$$((x - \frac{1}{n}, x] \times \{0\}) \cup ((x - \frac{1}{n}, x) \times \{1\}) \text{ if } x \in (0, 1] \text{ and } n \in \mathbb{N}$$

and

$$((x, x + \frac{1}{n}) \times \{0\}) \cup ([x, x + \frac{1}{n}) \times \{1\}) \text{ if } x \in [0, 1) \text{ and } n \in \mathbb{N}.$$

For a set $A \subseteq X$ we denote

$$A^+ = \{x \in [0, 1] : (x, 1) \in A\} \text{ and } A^- = \{x \in [0, 1] : (x, 0) \in A\}.$$

It is not hard to verify that for every open or closed set $A \subseteq X$ we have $|A^+ \Delta A^-| \leq \aleph_0$. It follows that $|B^+ \Delta B^-| \leq \aleph_0$ for any Borel measurable set $B \subseteq X$.

Let $E = X_0$. Since $E^+ = \emptyset$ and $E^- = (0, 1]$, the set E is non-measurable. We show that E is an ambiguously 1-embedded set. Indeed, let $A \subseteq E$ be an F_σ - and G_δ -subset of E . Then $B = E \setminus A$ is also an F_σ - and G_δ -subset of E . Let $\tilde{A}$ and $\tilde{B}$ be G_δ -sets in X such that $A = \tilde{A} \cap E$ and $B = \tilde{B} \cap E$. The inequalities $|\tilde{A}^+ \Delta \tilde{A}^-| \leq \aleph_0$ and $|\tilde{B}^+ \Delta \tilde{B}^-| \leq \aleph_0$ imply that $|C| \leq \aleph_0$, where $C = \tilde{A} \cap \tilde{B}$. Hence, C is an F_σ -set in X . Moreover, C is a G_δ -set in X . Therefore, $\tilde{A} \setminus C$ and $\tilde{B} \setminus C$ are G_δ -sets in X . According to Lemma 4.2, there is an F_σ - and G_δ -set D in X such that $\tilde{A} \setminus C \subseteq D$ and $D \cap (\tilde{B} \setminus C) = \emptyset$. Then $D \cap E = A$. $\square$

5. EXTENSION OF REAL-VALUED K_α -FUNCTIONS

Analogues of Proposition 5.1 and Theorem 5.3 for $\alpha = 1$ were proved in [7].

Proposition 5.1. *Let X be a topological space, $E \subseteq X$ and $0 < \alpha < \omega_1$. Then the following conditions are equivalent:*

- (i) E is K_α^* -embedded in X ;
- (ii) E is ambiguously α -embedded in X ;

(iii) $(X, E, [c, d])$ has the K_α -extension property for any segment $[c, d] \subseteq \mathbb{R}$.

Proof. (i) $\implies$ (ii). Take an arbitrary functionally ambiguous set A of the class α in E and consider its characteristic function χ_A . Then $\chi_A \in K_\alpha^*(E)$, as is easy to check. Let $f \in K_\alpha(X)$ be an extension of χ_A . Then the sets $f^{-1}(1)$ and $f^{-1}(0)$ are disjoint and belong to the α 'th functionally multiplicative class in X . According to Lemma 4.2 there exists a functionally ambiguous set B of the class α in X such that $f^{-1}(1) \subseteq B$ and $B \cap f^{-1}(0) = \emptyset$. It remains to notice that $B \cap E = f^{-1}(1) \cap E = \chi_A^{-1}(1) = A$. Hence, E is an ambiguously α -embedded set in X .

(ii) $\implies$ (iii). Let $f \in K_\alpha(E, [c, d])$. Define

$$h_1(x) = \begin{cases} f(x), & \text{if } x \in E, \\ \inf f(E), & \text{if } x \in X \setminus E, \end{cases}$$

$$h_2(x) = \begin{cases} f(x), & \text{if } x \in E, \\ \sup f(E), & \text{if } x \in X \setminus E, \end{cases}$$

Then $c \leq h_1(x) \leq h_2(x) \leq d$ for all $x \in X$.

We prove that for any reals $a < b$ there exists a function $h \in K_\alpha(X)$ such that

$$h_2^{-1}([c, a]) \subseteq h^{-1}(0) \quad \text{and} \quad h_1^{-1}([b, d]) \subseteq h^{-1}(1).$$

Fix $a < b$. Without loss of generality we may assume that

$$\inf f(E) \leq a < b \leq \sup f(E).$$

Denote

$$A_1 = f^{-1}([c, a]), \quad A_2 = f^{-1}([b, d]).$$

Then A_1 and A_2 are disjoint sets of the α 'th functionally multiplicative class in E . Using Lemma 4.2, we choose a functionally ambiguous set C of the class α in E such that $A_1 \subseteq C$ and $C \cap A_2 = \emptyset$. Since E is an ambiguously α -embedded set in X , there exists such a functionally ambiguous set D of the class α in X that $D \cap E = C$. Moreover, by Proposition 4.1 there exist sets B_1 and B_2 of the α 'th functionally multiplicative class in X such that $A_i = E \cap B_i$ when $i = 1, 2$. Let

$$\tilde{A}_1 = D \cap B_1, \quad \tilde{A}_2 = (X \setminus D) \cap B_2.$$

Then the sets $\tilde{A}_1$ and $\tilde{A}_2$ are disjoint and belong to the α 'th functionally multiplicative class in X . Moreover, $A_1 = E \cap \tilde{A}_1$ and $A_2 = E \cap \tilde{A}_2$. According to Proposition 3.4 there is a function $h \in K_\alpha^*(X)$ such that

$$h^{-1}(0) = \tilde{A}_1 \quad \text{and} \quad h^{-1}(1) = \tilde{A}_2.$$

According to [12, Theorem 3.2] there exists a function $g \in K_\alpha(X)$ such that

$$h_1(x) \leq g(x) \leq h_2(x)$$

for all $x \in X$. Clearly, g is an extension of f and $g \in K_\alpha(X, [c, d])$.

(iii) $\implies$ (i). Let $f \in K_\alpha^*(E)$ and let $|f(x)| \leq C$ for all $x \in E$. Consider a function $g \in K_\alpha(X)$ which is an extension of f . Define a function $r : \mathbb{R} \rightarrow [-C, C]$, $r(x) = \min\{C, \max\{x, -C\}\}$. Obviously, r is continuous. Let $h = r \circ g$. Then $h \in K_\alpha^*(X)$ and $h|_E = f$. Hence, E is K_α^* -embedded in X . $\square$

Lemma 5.2. *Let $0 < \alpha < \omega_1$, X be a topological space and let $E \subseteq X$ be such an α -embedded set in X that for any set A of the α 'th functionally multiplicative class in X such that $E \cap A = \emptyset$ the sets E and A are α -separated. Then E is an ambiguously α -embedded set.*

Proof. Consider a functionally ambiguous set C of the class α in E and denote $C_1 = C$, $C_2 = E \setminus C$. Then there exist sets $\tilde{C}_1$ and $\tilde{C}_2$ of the α 'th functionally multiplicative class in X such that $\tilde{C}_i \cap E = C_i$ when $i = 1, 2$. Then the set $A = \tilde{C}_1 \cap \tilde{C}_2$ is of the α 'th functionally multiplicative class in X and $A \cap E = \emptyset$. Let $h \in K_\alpha(X)$ be a function such that $E \subseteq h^{-1}(0)$ and $A \subseteq h^{-1}(1)$. Denote $H = h^{-1}(0)$ and $H_i = H \cap \tilde{C}_i$ when $i = 1, 2$. Since H_1 and H_2 are disjoint sets of the α 'th functionally multiplicative class in X , by Lemma 4.2 there is a functionally ambiguous set D of the class α in X such that $H_1 \subseteq D \subseteq X \setminus H_2$. Obviously, $D \cap E = C$. $\square$

Theorem 5.3. *Let $0 < \alpha < \omega_1$ and let E be a subset of a topological space X . Then the following conditions are equivalent:*

- (i) E is K_α -embedded in X ;
- (ii) E is α -embedded in X and for any set A of the α 'th functionally multiplicative class in X such that $E \cap A = \emptyset$ the sets E and A are α -separated.

Proof. (i) $\implies$ (ii). Let $C \subseteq E$ be a set of the α 'th functionally multiplicative class in E . Then by Lemma 3.3 we choose a function $f \in K_\alpha^*(E)$ such that $C = f^{-1}(0)$. If $g \in K_\alpha(X)$ is an extension of f , then the set $B = g^{-1}(0)$ belongs to the α 'th functionally multiplicative class in X and $B \cap E = C$. Hence, E is an α -embedded set in X .

Now consider a set A of the α 'th functionally multiplicative class in X such that $E \cap A = \emptyset$. According to Lemma 3.3 there is a function $h \in K_\alpha^*(X)$ such that $A = h^{-1}(0)$. For all $x \in E$ let $f(x) = \frac{1}{h(x)}$. Then $f \in K_\alpha(E)$. Let $g \in K_\alpha(X)$ be an extension of f . For all $x \in X$ let $\varphi(x) = g(x) \cdot h(x)$. Clearly, $\varphi \in K_\alpha(X)$. It is not hard to verify that $E \subseteq \varphi^{-1}(1)$ and $A \subseteq \varphi^{-1}(0)$.

(ii) $\implies$ (i). Remark that according to Lemma 5.2 the set E is ambiguously α -embedded in X .

Let $f \in K_\alpha(E)$ and let $\varphi : \mathbb{R} \rightarrow (-1, 1)$ be a homeomorphism. Using Proposition 5.1 to the function $\varphi \circ f : E \rightarrow [-1, 1]$ we have that there exists a function $h \in K_\alpha(X, [-1, 1])$ such that $h|_E = \varphi \circ f$. Let

$$A = h^{-1}(-1) \cup h^{-1}(1).$$

Then A belongs to the α 'th functionally multiplicative class in X and $A \cap E = \emptyset$. Therefore, there exists a function $\psi \in K_\alpha(X)$ such that $A \subseteq \psi^{-1}(0)$ and $E \subseteq \psi^{-1}(1)$. For all $x \in X$ define

$$g(x) = \varphi^{-1}(h(x) \cdot \psi(x)).$$

Remark that $g \in K_\alpha(X)$ and $g|_E = f$. $\square$

Corollary 5.4. *Let $0 < \alpha < \omega_1$ and let E be a subset of the α 'th functionally multiplicative class of a topological space X . Then the following conditions are equivalent:*

- (i) E is K_α -embedded in X ;
- (ii) E is α -embedded in X .

6. K_1^* -EMBEDDING VERSUS K_1 -EMBEDDING

A family $\mathcal{U}$ of non-empty open sets of a space X is called a π -base [4] if for any non-empty open set V of X there is $U \in \mathcal{U}$ with $V \subseteq U$.

Proposition 6.1. *Let X be a perfect space of the first category with a countable π -base. Then there exist disjoint F_σ - and G_δ -subsets A and B of X which are dense in X and $X = A \cup B$.*

Proof. Let $(V_n : n \in \mathbb{N})$ be a π -base in X and $X = \bigcup_{n=1}^{\infty} X_n$, where X_n is a closed nowhere dense subset of X for every $n \geq 1$. Let $E_1 = X_1$ and $E_n = X_n \setminus \bigcup_{k < n} X_k$ for $n \geq 2$. Then E_n is a nowhere dense F_σ - and G_δ -subset of X for every $n \geq 1$, $E_n \cap E_m = \emptyset$ if $n \neq m$, and $X = \bigcup_{n=1}^{\infty} E_n$.

Let $m_0 = 0$. We choose a number $n_1 \geq 1$ such that $(\bigcup_{n=1}^{n_1} E_n) \cap V_1 \neq \emptyset$ and let $A_1 = \bigcup_{n=1}^{n_1} E_n$. Since $\overline{X \setminus A_1} = X$, there exists a number $m_1 > n_1$ such that $(\bigcup_{n=n_1+1}^{m_1} E_n) \cap V_1 \neq \emptyset$. Set $B_1 = \bigcup_{n=n_1+1}^{m_1} E_n$. It follows from the equality $\overline{X \setminus (A_1 \cup B_1)} = X$ that there exists $n_2 > m_1$ such that $(\bigcup_{n=m_1+1}^{n_2} E_n) \cap V_2 \neq \emptyset$. Further, there is such $m_2 > n_2$ that $(\bigcup_{n=n_2+1}^{m_2} E_n) \cap V_2 \neq \emptyset$. Let $A_2 = \bigcup_{n=m_1+1}^{n_2} E_n$ and $B_2 = \bigcup_{n=n_2+1}^{m_2} E_n$. Repeating this process, we obtain the sequence of numbers

$$m_0 < n_1 < m_1 < \dots < n_k < m_k < n_{k+1} < \dots$$

and the sequence of sets

$$A_k = \bigcup_{n=m_{k-1}+1}^{n_k} E_n, \quad B_k = \bigcup_{n=n_k+1}^{m_k} E_n, \quad k \geq 1,$$

such that $A_k \cap V_k \neq \emptyset$ and $B_k \cap V_k \neq \emptyset$ for every $k \geq 1$.

Let $A = \bigcup_{k=1}^{\infty} A_k$ and $B = \bigcup_{k=1}^{\infty} B_k$. Clearly, $X = A \cup B$, $A \cap B = \emptyset$ and $\overline{A} = \overline{B} = X$. Moreover, A and B are F_σ -sets in X . Therefore, A and B are F_σ - and G_δ -subsets of X . $\square$

We say that a topological space X *hereditarily has a countable π -base* if every its closed subspace has a countable π -base.

Proposition 6.2. *Let X be a hereditarily Baire space, E be a perfectly normal ambiguously 1-embedded subspace of X which hereditarily has a countable π -base. Then E is a hereditarily Baire space.*

Proof. Assume that E is not a hereditarily Baire space. Then there exists a nonempty closed set $C \subseteq X$ of the first category. Notice that C is a perfectly normal space with a countable π -base. According to Proposition 6.1 there exist disjoint dense F_σ - and G_δ -subsets A and B of C such that $C = A \cup B$. Since C is F_σ - and G_δ -set in E , the sets A and B are also F_σ and G_δ in E . Therefore there exist disjoint functionally F_σ - and G_δ -subsets $\tilde{A}$ and $\tilde{B}$ of X such that $A = \tilde{A} \cap E$ and $B = \tilde{B} \cap E$. Notice that the sets $\tilde{A}$ and $\tilde{B}$ are dense in $\overline{C}$. Taking into account that X is hereditarily Baire, we have that $\overline{C}$ is a Baire space. It follows a contradiction, since $\tilde{A}$ and $\tilde{B}$ are disjoint dense G_δ -subsets of $\overline{C}$. $\square$

Remark that there exist a metrizable separable Baire space X and its ambiguously 1-embedded subspace E which is not a Baire space. Indeed, let $X = (\mathbb{Q} \times \{0\}) \cup (\mathbb{R} \times (0, 1])$ and $E = \mathbb{Q} \times \{0\}$. Then E is closed in X . Therefore, any F_σ - and G_δ -subset C of E is also F_σ - and G_δ - in X . Hence, E is an ambiguously 1-embedded set in X .

Theorem 6.3. *Let X be a hereditarily Baire space and let $E \subseteq X$ be its perfect Lindelöf subspace which hereditarily has a countable π -base. Then E is K_1^* -embedded in X if and only if E is K_1 -embedded in X .*

Proof. Since the sufficiency is obvious, we only need to prove the necessity.

According to Proposition 5.1 the set E is ambiguously 1-embedded in X . Using Proposition 6.2, we have E is a hereditarily Baire space. Since E is Lindelöf, Proposition 3.9 (b) implies that E is 1-separated from any functionally G_δ -set A of X such that $A \cap E = \emptyset$. Therefore, by Theorem 5.3 the set E is K_1 -embedded in X . $\square$

7. A GENERALIZATION OF THE KURATOWSKI THEOREM

K. Kuratowski [11, p. 445] proved that every mapping $f \in K_\alpha(E, Y)$ has an extension $g \in K_\alpha(X, Y)$ if the case X is a metric space, Y is a Polish space and $E \subseteq X$ is a set of the multiplicative class $\alpha > 0$.

In this section we will prove that the Kuratowski Extension Theorem is still valid if X is a topological space and E is a K_α -embedded subset of X .

We say that a subset A of a space X is *discrete* if any point $a \in A$ has a neighborhood $U \subseteq X$ such that $U \cap A = \{a\}$.

Theorem 7.1 (Theorem 2.11 [8]). *Let X be a topological space, Y be a metrizable separable space, $0 \leq \alpha < \omega_1$ and $f \in K_\alpha(X, Y)$. Then there exists a sequence $(f_n)_{n=1}^\infty$ such that*

- (i) $f_n \in K_\alpha(X, Y)$ for every n ;

- (ii) $(f_n)_{n=1}^\infty$ is uniformly convergent to f ;
- (iii) $f_n(X)$ is at most countable and discrete for every n .

Proof. Consider a metric d on Y which generates its topological structure. Since (Y, d) is metric separable space, for every n there is a subset $Y_n = \{y_{i,n} : i \in I_n\}$ of Y such that Y_n is discrete, $|I_n| \leq \aleph_0$ and for any $y \in Y$ there exists $i \in I_n$ such that $d(y, y_{i,n}) < 1/n$ (see [11, p. 226]).

For every $n \in \mathbb{N}$ and $i \in I_n$ put $A_{i,n} = \{x \in X : d(f(x), y_{i,n}) < 1/n\}$. Then each $A_{i,n}$ belongs to the α 'th functionally additive class in X and $\bigcup_{i \in I_n} A_{i,n} = X$ for every n . According to Lemma 3.2 for every n we can choose a sequence $(F_{i,n})_{i \in I_n}$ of disjoint functionally ambiguous sets of the class α such that $F_{i,n} \subseteq A_{i,n}$ and $\bigcup_{i \in I_n} F_{i,n} = X$.

For all $x \in X$ and $n \in \mathbb{N}$ let $f_n(x) = y_{i,n}$ if $x \in F_{i,n}$ for some $i \in I_n$. Notice that $f_n \in K_\alpha(X, Y)$ for every $n \in \mathbb{N}$.

It remains to prove that the sequence $(f_n)_{n=1}^\infty$ is uniformly convergent to f . Indeed, fix $x \in X$ and $n \in \mathbb{N}$. Then there exists $i \in I_n$ such that $x \in F_{i,n}$. Since $F_{i,n} \subseteq A_{i,n}$, $d(f(x), f_n(x)) = d(f(x), y_{i,n}) < \frac{1}{n}$, which completes the proof. $\square$

Recall that a family $(A_s : s \in S)$ of subsets of a topological space X is called a *partition* of X if $X = \bigcup_{s \in S} A_s$ and $A_s \cap A_t = \emptyset$ for all $s \neq t$.

Proposition 7.2. *Let $0 < \alpha < \omega_1$, X be a topological space, $E \subseteq X$ be an α -embedded set which is α -separated from any disjoint with it set of the α 'th functionally multiplicative class in X and let $(A_n : n \in \mathbb{N})$ be a partition of E by functionally ambiguous sets of the class α in E . Then there is a partition $(B_n : n \in \mathbb{N})$ of X by functionally ambiguous sets of the class α in X such that $A_n = E \cap B_n$ for every $n \in \mathbb{N}$.*

Proof. According to Proposition 5.2 for every $n \in \mathbb{N}$ there exists a functionally ambiguous set D_n of the class α in X such that $A_n = D_n \cap E$. By the assumption there exists a function $f \in K_\alpha(X)$ such that $E \subseteq f^{-1}(0)$ and $X \setminus \bigcup_{n=1}^\infty D_n \subseteq f^{-1}(1)$.

Let $D = f^{-1}(0)$. Then the set $X \setminus D$ is of the α 'th functionally additive class in X . Then there exists a sequence $(E_n)_{n=1}^\infty$ of functionally ambiguous set of the class α in X such that $X \setminus D = \bigcup_{n=1}^\infty E_n$. For every $n \in \mathbb{N}$ denote $C_n = E_n \cup D_n$. Then

all the sets C_n are functionally ambiguous of the class α in X and $\bigcup_{n=1}^\infty C_n = X$.

Let $B_1 = C_1$ and $B_n = C_n \setminus \left(\bigcup_{k < n} C_k \right)$ for $n \geq 2$. Clearly, every B_n is a functionally ambiguous set of the class α in X , $B_n \cap B_m = \emptyset$ if $n \neq m$ and $\bigcup_{n=1}^\infty B_n = \bigcup_{n=1}^\infty C_n = X$. Moreover, $B_n \cap E = A_n$ for every $n \in \mathbb{N}$. $\square$

Let $0 \leq \alpha < \omega_1$, X and Y be topological spaces and $E \subseteq X$. We say that a collection (X, E, Y) has the K_α -extension property if every mapping $f \in K_\alpha(E, Y)$ can be extended to a mapping $g \in K_\alpha(X, Y)$.

Theorem 7.3. *Let $0 < \alpha < \omega_1$ and let E be a subset of a topological space X . Then the following conditions are equivalent:*

- (i) E is K_α -embedded in X ;
- (ii) (X, E, Y) has the K_α -extension property for any Polish space Y .

Proof. Since the implication (ii) $\Rightarrow$ (i) is obvious, we only need to prove the implication (i) $\Rightarrow$ (ii). Let Y be a Polish space with a metric d which generates its topological structure such that (Y, d) is complete and let $f \in K_\alpha(E, Y)$.

It follows from Theorem 7.1 that there exists a sequence of mappings $f_n \in K_\alpha(E, Y)$ which is uniformly convergent to f on E . Moreover, for every $n \in \mathbb{N}$ the set $f_n(E) = \{y_{i_n, n} : i_n \in I_n\}$ is at most countable and discrete. We may assume that each $f_n(E)$ consists of distinct points.

For every $n \in \mathbb{N}$ and for each $(i_1, \dots, i_n) \in I_1 \times \dots \times I_n$ let

$$B_{i_1, \dots, i_n} = f_1^{-1}(y_{i_1, 1}) \cap \dots \cap f_n^{-1}(y_{i_n, n}).$$

Then for each $i_1 \in I_1, \dots, i_n \in I_n$ the set $B_{i_1, \dots, i_n}$ is functionally ambiguous of the class α in E and the family $(B_{i_1, \dots, i_n} : i_1 \in I_1, \dots, i_n \in I_n)$ is a partition of E for every $n \in \mathbb{N}$. By Proposition 7.2 we choose a sequence of systems of functionally ambiguous sets $D_{i_1, \dots, i_n}$ of the class α in X such that

- (1) $D_{i_1, \dots, i_n} \cap E = B_{i_1, \dots, i_n}$ for every $n \in \mathbb{N}$ and $(i_1, \dots, i_n) \in I_1 \times \dots \times I_n$;
- (2) $(D_{i_1, \dots, i_n} : i_1 \in I_1, \dots, i_n \in I_n)$ is a partition of X for every $n \in \mathbb{N}$.

For all $n \in \mathbb{N}$ and $(i_1, \dots, i_n) \in I_1 \times \dots \times I_n$ let

- (3) $D_{i_1, \dots, i_n} = \emptyset$, if $B_{i_1, \dots, i_n} = \emptyset$.

Notice that the system $(B_{i_1, \dots, i_n, i_{n+1}} : i_{n+1} \in I_{n+1})$ forms a partition of the set $B_{i_1, \dots, i_n}$ for every $n \in \mathbb{N}$.

For all $i_1 \in I_1$ let

$$C_{i_1} = D_{i_1}.$$

Assume that for some $n \geq 1$ the system $(C_{i_1, \dots, i_n} : i_1 \in I_1, \dots, i_n \in I_n)$ of functionally ambiguous sets of the class α in X is already defined and

- (A) $B_{i_1, \dots, i_n} = E \cap C_{i_1, \dots, i_n}$;
- (B) $(C_{i_1, \dots, i_n} : i_1 \in I_1, \dots, i_n \in I_n)$ is a partition of X ;
- (C) $C_{i_1, \dots, i_n} = \emptyset$ if $B_{i_1, \dots, i_n} = \emptyset$;
- (D) $(C_{i_1, \dots, i_{n-1}, i_n} : i_n \in I_n)$ is a partition of the set $C_{i_1, \dots, i_{n-1}}$.

Fix $i_1, \dots, i_n$. Since the set $K = C_{i_1, \dots, i_n} \setminus \bigcup_{k \in I_{n+1}} D_{i_1, \dots, i_n, k}$ is of the α 'th functionally

multiplicative class in X and $K \cap E = \emptyset$, there exists a set H of the α 'th functionally multiplicative class in X such that $E \subseteq H \subseteq X \setminus K$. Using [8, Lemma 2.1] we obtain that there exists a sequence $(A_k)_{k=1}^\infty$ of disjoint functionally

ambiguous sets of the class α in X such that

$$C_{i_1, \dots, i_n} \setminus H = \bigcup_{k=1}^{\infty} A_k.$$

Let

$$M_{i_1, \dots, i_n, i_{n+1}} = \emptyset, \quad \text{if } D_{i_1, \dots, i_n, i_{n+1}} = \emptyset,$$

and

$$M_{i_1, \dots, i_n, i_{n+1}} = (A_{i_{n+1}} \cup D_{i_1, \dots, i_n, i_{n+1}}) \cap C_{i_1, \dots, i_n}, \quad \text{if } D_{i_1, \dots, i_n, i_{n+1}} \neq \emptyset.$$

Now let

$$C_{i_1, \dots, i_n, 1} = M_{i_1, \dots, i_n, 1},$$

and

$$C_{i_1, \dots, i_n, i_{n+1}} = M_{i_1, \dots, i_n, i_{n+1}} \setminus \bigcup_{k < i_{n+1}} M_{i_1, \dots, i_n, k} \quad \text{if } i_{n+1} > 1.$$

Then for every $n \in \mathbb{N}$ the system $(C_{i_1, \dots, i_n} : i_1 \in I_1, \dots, i_n \in I_n)$ of functionally ambiguous sets of the class α in X has the properties (A)–(D).

For each $n \in \mathbb{N}$ and $x \in X$ let

$$g_n(x) = y_{i_n, n},$$

if $x \in C_{i_1, \dots, i_n}$. It is not hard to prove that $g_n \in K_\alpha(X, Y)$.

We show that the sequence $(g_n)_{n=1}^\infty$ is uniformly convergent on X . Indeed, let $x_0 \in X$ and $n, m \in \mathbb{N}$. Without loss of generality, we may assume that $n \geq m$. By the property (B), $x_0 \in C_{i_1, \dots, i_n} \cap C_{j_1, \dots, j_m}$. It follows from (B) and (D) that $i_1 = j_1, \dots, i_m = j_m$. Take an arbitrary point x from the set $B_{i_1, \dots, i_n}$, the existence of which is guaranteed by the property (C). Then $f_m(x) = y_{i_m, m} = g_m(x_0)$ and $f_n(x) = y_{i_n, n} = g_n(x_0)$. Since the sequence $(f_n)_{n=1}^\infty$ is uniformly convergent on E , $\lim_{n, m \rightarrow \infty} d(y_{i_m, m}, y_{i_n, n}) = 0$. Hence, the sequence $(g_n)_{n=1}^\infty$ is uniformly convergent on X .

Since Y is a complete space, for all $x \in X$ define $g(x) = \lim_{n \rightarrow \infty} g_n(x)$. According to the property (A), $g(x) = f(x)$ for all $x \in E$. Moreover, $g \in K_\alpha(X, Y)$ as a uniform limit of functions from the class K_α . $\square$

8. OPEN PROBLEMS

Question 8.1. *Does there exist a completely regular not perfectly normal space in which any functionally G_δ -set is 1-embedded?*

Question 8.2. *Does there exist a completely regular not perfectly normal space in which any set is 1-embedded?*

Question 8.3. *Do there exist a normal space and its functionally G_δ -subset which is not 1-embedded?*

Question 8.4. *Do there exist a topological space X and its subspace E such that E is K_1^* -embedded and is not K_1 -embedded in X ?*

9. ACKNOWLEDGMENTS

The author would like to thank the referee for his helpful and constructive comments that greatly contributed to improving the final version of the paper.

REFERENCES

- [1] Blair R. *Filter characterization of z -, C^* -, and C -embeddings*, Fund. Math. **90** (1976), 285–300.
- [2] Blair R., Hager A. *Extensions of zero-sets and of real-valued functions*, Math. Zeit. **136** (1974), 41–52.
- [3] Corson H. *Normality in subsets of product spaces*, Amer. J. Math. **81** (1959), 785–796.
- [4] *Encyclopedia of General Topology*. Edited by Klaas Pieter Hart, Jun-iti Nagata and Jerry E. Vaughan, Elsevier (2004).
- [5] Engelking R. *General Topology. Revised and completed edition*. Heldermann Verlag, Berlin (1989).
- [6] Gillman L., Jerison M. *Rings of continuous functions*, Van Nostrand, Princeton (1960).
- [7] Kalenda, O., Spurný, J. *Extending Baire-one functions on topological spaces*, Topol. Appl. **149** (2005), 195–216.
- [8] Karlova O. *Baire classification of mappings which are continuous with respect to the first variable and of the α 'th functionally class with respect to the second variable*, Mathematical Bulletin NTSH, **2** (2005), 98–114 (in Ukrainian).
- [9] Karlova O. *Classification of separately continuous functions with values in σ -metrizable spaces*, Applied General Topology **13** (2) (2012), 167–178.
- [10] Kombarov A., Malykhin V. *On Σ -products*, DAN SSSR, **213** (1973), 774–776 (in Russian).
- [11] Kuratowski K. *Topology, V.1*, Moscow: Mir (1966) (in Russian).
- [12] Lukeš J., Malý J., Zajíček L. *Fine Topology Methods in Real Analysis and Potentail Theory*, Springer-Verlag, (1986).
- [13] Ohta H. *Extension properties and the Niemytski plane*, Applied General Topology, **1** (1) (2000), 45–60.

CHERNIVTSI NATIONAL UNIVERSITY, DEPARTMENT OF MATHEMATICAL ANALYSIS, KOT-SJUBYNS'KOHO 2, CHERNIVTSI 58012, UKRAINE